

***Mylabris aschnae* sp. nov., a new blister beetle from Mahallat, Markazi Province, Iran (Coleoptera: Meloidae)**

by Dewanand Makhan* & Somayeh Ezzatpanah **

*Willem Bilderdijkhove 19, 3438 PM Nieuwegein, The Netherlands (email: dewanandmakhan@yahoo.co.uk)

**School of Biology, College of Science, University of Tehran, Tehran, Iran (email: s.ezzatpanah@gmail.com)

Makhan, D. & Ezzatpanah, S. (2011). *Mylabris aschnae* sp. nov., a new blister beetle from Mahallat, Markazi Province, Iran (Coleoptera: Meloidae). *Calodema*, 192: 1-6.

Abstract: A new *Mylabris* species of Meloidae (Coleoptera) is described from Mahallat, Markazi Province, Iran.

Key words: *Mylabris*, Meloidae, Coleoptera, Iran.

Introduction

Nikbakhtzadeh & Tigrari (2002 2007) provided only the habitus of the four species of *Mylabris* Fabricius, 1775 (Coleoptera: Meloidae) and the figures of the aedeagus of each are absent. The species are *Mylabris guerini* Chevrolat, 1840, *Mylabris impressa* (Chevrolat, 1837), *Mylabris schreibersi* Reiche 1865 and *Mylabris variabilis* (Pallas 1781). This material was not available to us and further study of these is necessary. It should be noted that the claws of the genus *Mylabris* Fabricius 1775 possess teeth in some species while in others the claws lack teeth.

Mylabris aschnae Makhan & Ezzatpanah, sp. nov. appears to be allied to *Mylabris (Eumylabris) impressa* (Chevrolat, 1837) in the papers by Nikbakhtzade & Tigrari (2002, 2007), but *Mylabris (Eumylabris) impressa* (Chevrolat, 1837) is absent from Iran. *Mylabris (Eumylabris) impressa* (Chevrolat, 1837) from Spain has 6 rounded spots on each elytron.

The specimens were collected from flowers of *Citrullus vulgaris* L. (Cucurbitaceae).

The holotype will be deposited in the collection of the School of Biology, College of Science, University of Tehran, Tehran, Iran. Paratypes are deposited in the first author's collection.

Taxonomic description

Mylabris aschnae Makhan & Ezzatpanah, sp. nov. (Figs. 1-9)

Holotype: Male. Iran, Markazi Province, Mahallat, 1.7.2011, D. Makhan & S. Ezzatpanah. Paratypes: 55 specimens.

Holotype description: Male. Length 24.5 mm, width 6.0 mm. Head black, length 6.0 mm, width 4.0 mm. Eyes large, maxilla palpus black, with fine punctures, with black setae. Antennae black, 11-segmented, antennomeres 1-2, 4-11 longer than wide, antennomere 3 wider than long, with black setae. Pronotum black, length 4.0 mm, width 3.5 mm, widest at base, with coarse punctures, with a large shallow depression at middle and with a large shallow depression at base, with black setae. Elytra length 17.0 mm, width 6.0 mm. Elytron

orange, with two black transverse fasciae and two rounded black spots at anterior side, with fine punctures and with black setae. Legs: Figs 7-8, black, with black setae, tarsus with bristles and the claws with teeth. Male aedeagus: Fig. 6, with two teeth and with a hook at apex.

Etymology: This species is named after Dewanand's daughter, Aschna Makhan.

Remarks: *Mylabris aschneae* Makhan & Ezzatpanah, sp. nov. appears to be most closely allied to *Mylabris calida* (Pallas, 1782) an Turano-Mediterranean species, but there is a difference in the shape of the male aedeagus. The aedeagus of *Mylabris (Eumylabris) calida* (Pallas, 1782) is wide at the apical third and the male aedeagus of *Mylabris aschneae* Makhan & Ezzatpanah, sp. nov. is narrow.

References

- Nikbakhtzadeh, M.R. & Tigrari, S. (2002). Blister Beetles (Coleoptera: Meloidae) in Nahavand County (Hamedan Province, Iran) and their ecological relationship to other Coleopteran families. *Iranian Journal of Public Health*, 31 (1-2): 55-62.
Nikbakhtzadeh, M.R. & Tigrari , S. (2007). Medically important beetles (Insecta: Coleoptera) of Iran. *Journal of Venom and Animal Toxins including Tropical Diseases*, 14 (4): 597-618.

Acknowledgements

Thanks are expressed to Trevor J. Hawkeswood of Sydney, Australia for editorial assistance, Zhao Pan & Guo-Dong Ren from the College of Life Sciences, Hubei University, Baoding, Hebei, China and Marco A. Bologna Dipartimento di Biologia ambientale, Università Roma Tre, Viale Marconi, Roma, Italy, also for all various assistances.



Fig. 1. *Mylabris aschneae* Makhan & Ezzatpanah, sp. nov., habitus, holotype, lateral view
(scale line = 4.0 mm).



Fig. 2. *Mylabris aschneae* Makhan & Ezzatpanah, sp. nov., habitus, paratype, dorsal view
(scale line = 5.0 mm).




Fig. 3. *Mylabris aschneae* Makhan & Ezzatpanah, sp. nov., holotype male speculum gastrale, dorsal view (scale line = 1.2 mm).

Fig. 4. *Mylabris aschneae* Makhan & Ezzatpanah, sp. nov., holotype, gonoforceps, lateral view (scale line = 1.0 mm).

Fig. 5. *Mylabris aschneae* Makhan & Ezzatpanah, sp. nov., holotype male, gonoforceps, dorsal view (scale line = 1.0 mm).




Fig. 6. *Mylabris aschneae* Makhan & Ezzatpanah, sp. nov., holotype male aedeagus, lateral view
(scale line = 1.5 mm).




Fig. 7. *Mylabris aschneae* Makhan & Ezzatpanah, sp. nov., holotype, claws, dorsal view
(scale line = 0.4 mm).




Fig. 8. *Mylabris aschneae* Makhan & Ezzatpanah, sp. nov., paratype male, right fore leg, ventral view
(scale line = 0.5 mm).




Fig. 9. *Mylabris aschneae* Makhan & Ezzatpanah, sp. nov., habitus, holotype, 9th tergite
(scale line = 0.5 mm).

Date of publication: 24 November 2011
Copyright: Dewanand Makhan & Somayeh Ezzatpanah
Editor: Dr T.J. Hawkeswood (www.calodema.com)
(Published as hard paper copy edition as well as electronic pdf)